



King Saud University  
**Journal of Saudi Chemical Society**

[www.ksu.edu.sa](http://www.ksu.edu.sa)  
[www.sciencedirect.com](http://www.sciencedirect.com)

**ORIGINAL ARTICLE**

# The combined role of heterogeneous catalysis and ultrasonic waves on the facile synthesis of 2,3-dihydroquinazolin-4(1*H*)-ones



Javad Safari <sup>\*</sup>, Soheila Gandomi-Ravandi

Laboratory of Organic Compound Research, Department of Organic Chemistry, College of Chemistry, University of Kashan, P.O. Box 87317-51167, Kashan, Islamic Republic of Iran

Received 9 December 2013; revised 18 April 2014; accepted 26 April 2014  
Available online 15 May 2014

**KEYWORDS**

Dihydroquinazolinone;  
Multi-component reaction;  
Isatoic anhydride;  
Supported catalyst

**Abstract** Pt–MWCNTs nanocomposites are found to be an excellent and efficient catalyst to promote one-pot and three-component coupling reaction of isatoic anhydride, aldehyde and ammonium acetate or primary aromatic amine to produce 2,3-dihydroquinazolin-4(1*H*)-one derivatives under ultrasound irradiation. This novel method has the advantages such as short reaction times, convenient manipulation, excellent yields and the use of effective catalyst.

© 2014 King Saud University. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

**1. Introduction**

Since the first multi-component reaction was reported in 1850 by Strecker [1], this methodology has emerged as an especially attractive synthetic strategy for rapid and efficient synthesis. Since the products are formed in a single synthetic operation, the structural diversity can be achieved simply by varying the reaction components which shows high atom economy and high selectivity [2,3]. 2,3-Dihydro-4(1*H*)-quinazolinones are important bicyclic heterocycles which have emerged as versatile biologically active compounds possessing applications as diuretic [4], antitumor [5], antibacterial [6], anticonvulsant [7]

and antihypertensive agents [8]. In addition, these compounds can easily be oxidized to their quinazolin-4(3*H*)-one analogues [9–11], which also are important pharmacologically active compounds [12–17]. Due to the significant interest in 2,3-dihydro-4(1*H*)-quinazolinones, several synthetic protocols have been developed for these heterocyclic compounds. The usual procedure involves one pot three-component reaction of isatoic anhydride, aldehydes and amines to synthesize 2,3-dihydro-4(1*H*)-quinazolinones. The reported catalysts for this one-pot procedure are included montmorillonite K-10 [18,19], silica sulfuric acid (SSA) [20,21], zinc(II) perfluorooctanoate [Zn(PFO)<sub>2</sub>] [22], MCM-41-SO<sub>3</sub>H [23], *p*-toluenesulfonic acid-paraformaldehyde copolymer (copolymer-PTSA) [24], Amberlyst-15 [25], gallium(III) triflate (Ga(OTf)<sub>3</sub>) [26], KAl(SO<sub>4</sub>)<sub>2</sub>·12H<sub>2</sub>O (alum) [27], molecular iodine (I<sub>2</sub>) [28–30], *p*-toluenesulfonic acid (*p*-TsOH) [31], aluminium tris(dihydrogen phosphate) (Al(H<sub>2</sub>PO<sub>4</sub>)<sub>3</sub>) [32], silica-bonded *N*-propylsulfamic acid (SBNPSA) [33], acetic acid [34], SiO<sub>2</sub>–FeCl<sub>3</sub> [35], SnCl<sub>2</sub> [36], ceric ammonium nitrate (CAN) [37], ionic liquids [38–40], silica-bonded S-sulfonic acid [41] and Al/Al<sub>2</sub>O<sub>3</sub> and Fe<sub>3</sub>O<sub>4</sub> nanoparticles [42,43]. However, some reported

<sup>\*</sup> Corresponding author. Tel.: +98 (0)361 591 2320; fax: +98 (0)361 591 2397.

E-mail address: [Safari@kashanu.ac.ir](mailto:Safari@kashanu.ac.ir) (J. Safari).

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier

methods have certain disadvantages such as tedious work-up, high reaction temperature and low yields. Therefore, development of a simple and efficient procedure is an important goal in the synthesis of 2,3-dihydroquinazolin-4(1*H*)-ones.

On the other hand, transition metal nanoparticles as exclusive materials have wide application in the homogeneous and heterogeneous catalysts. Among metal nanoparticles, Pt nanoparticles (NPs) have attracted significant attention due to their high catalytic activity [44]. It is well known that the specific electrocatalytic activity of platinum nanoparticles is strongly related to the size and dispersion of the particles, the supporting materials, preparation methods and their surface conditions. Highly distributed nanoscale Pt particles with nanometer size particles and narrow size distribution are ideal for high electrocatalytic activity owing to their large surface-to-volume ratio [45–53]. In fact, the choice of a suitable supporting material is an important aspect that may affect the performance of supported electrocatalysts due to interactions and surface reactivity [54,55]. Many researchers have demonstrated that, multi-walled carbon nanotubes (MWCNTs) are attractive supporting materials among the possible supports for Pt NPs catalyst, which can support a high dispersion of Pt nanoparticles owing to their particular morphology and large surface area [56–58]. As a result, several research papers reported that Pt NPs supported on CNTs increase the electrocatalytic activity [59,60] and also electrochemical stability [61,62]. In fact, Pt nanoparticles on the external walls contact easily with the reactant. Accordingly, various approaches have been devised to load platinum nanoparticles onto the surface of CNTs with a high specific surface area, superior catalytic activity and lower catalyst loading. These include wet chemical impregnation [63–65], electrode position [51,66–68], colloidal methods [58,69,70] and ion exchange [71,72].

Ultrasonic irradiation has been introduced as an eco-environmental technology in green chemistry to improve a large number of organic reactions [73,74]. Ultrasonic energy provides an unusual mechanism to generate high-energy chemistry owing to the extraordinary temperature and pressure generated by the cavitation bubble collapse [75]. The ‘hot-spot’ theory suggests that temperatures up to 5000 K and pressures up to 1000 bar are produced during this collapse [76–78]. The vibration movement of the ultrasonic waves causes growth of cavitation bubbles through the diffusion of solute vapor in the volume of these bubbles. After the growth process, the bubble collapse breaks the chemical bonds of the solute molecules [79]. Moreover, collapsed bubbles can carry smaller particles causing shocks among them and possibly causing surface amorphization which may result in their sintering [80].

To our knowledge, no studies have been reported on the preparation of quinazolinones using ultrasonic energy in the presence of Pt–MWCNTs nanocomposites. Herein, we have developed a novel methodology to prepare 2,3-dihydroquinazolin-4(1*H*)-ones by employing isatoic anhydride, aldehydes and ammonium acetate or primary aromatic amines using Pt–MWCNTs as an efficient catalyst under sonication conditions (Scheme 1).

## 2. Experimental

All materials were from commercial reagent grade and were used as received. MWCNTs were purchased from Nanotech


Port Co. (Taiwan). These MWCNTs were produced via the chemical vapor deposition (CVD) method. The outer diameter of MWCNTs was between 10 and 20 nm. Melting points were obtained in open capillary tubes and were measured on an Electrothermal MK3 apparatus and were uncorrected.  $^1\text{H}$  NMR and  $^{13}\text{C}$  NMR spectra were measured with a Bruker DRX-400 spectrometer using tetramethylsilane (TMS) as an internal standard and DMSO- $d_6$  as the solvent at room temperature. Chemical shifts ( $\delta$ ) are reported in ppm downfield from TMS and coupling constant ( $J$ ) values are given in Hz. FT-IR spectra were determined on a Perkin Elmer FT-IR 550 spectrophotometer using the pressed KBr. The progress of the reactions was monitored by thin-layer chromatography (TLC), which was visualized with UV light. The UV spectra were recorded on a UV–Vis Varian Perkin–Elmer UV 550-S spectrophotometer in chloroform. X-ray powder diffraction (XRD) experiment was carried out in a Holland Philips Xpert X-ray powder diffraction (XRD) diffractometer (CuK $\alpha$ , radiation,  $\lambda = 0.154056$  nm), at a scanning speed of  $2^\circ/\text{min}$  from  $10^\circ$  to  $100^\circ$  ( $2\theta$ ). The prepared Pt–MWCNTs nanocomposites were also characterized by a TESCAN model VEGA II scanning electron microscope (SEM) operated at a 15 kV accelerating voltage. The Raman spectra were measured on a Bruker SENTERRA spectrometer with spectral range: 200–3500  $\text{cm}^{-1}$  and Laser wavenumber 785 nm. Products were characterized by comparison of their physical and spectral data with those of authentic samples.

### 2.1. Synthesis of nanocatalyst


Pt–MWCNTs were prepared according to mentioned procedure in literature [81]. Briefly, the MWCNTs were purified in 2.5 M  $\text{HNO}_3$  and 0.5 M  $\text{H}_2\text{SO}_4$  at  $60^\circ\text{C}$  under sonication for 6 h. Then, modified MWCNTs were washed with distilled water to neutral pH and were dried in air at  $100^\circ\text{C}$  for 1 h. Then, the modified MWCNTs were added to 0.004 M  $\text{PtCl}_4$  aqueous solution and the suspension was sonicated at room temperature for 30 min with the pH value of the solution adjusted to 8–10 by adding NaOH solution. An excess of  $\text{NaBH}_4$  was added as reductant to reduce Pt. Then, the pH was maintained above 7. The reaction mixture was sonicated for 10 min and then was left to stand at room temperature for 24 h. Next, the product was filtered and was washed with distilled water and was dried in air at room temperature (Fig. 1).

### 2.2. General procedure for the synthesis of 2,3-dihydroquinazolin-4(1*H*)-ones under ultrasound irradiation

Pt–MWCNTs (0.04 g) were added to a mixture of isatoic anhydride (1.00 mmol, 0.13 g), aniline (1.20 mmol) or ammonium acetate (1.10 mmol) and aromatic aldehyde (1.00 mmol) in ethanol (5.00 mL). Then, the reaction mixture was irradiated in an ultrasound bath at temperature of  $60^\circ\text{C}$  for appropriate time. After completion of the reaction as indicated by TLC (petroleum ether–EtOAc, 7:3), the reaction mixture was cooled to room temperature, was diluted with hot ethanol and the solid catalyst was separated by filtration.  $\text{H}_2\text{O}$  (5.00 mL) was added and the precipitated product was filtered and was recrystallized from ethanol.



**Scheme 1** Ultrasonic activated synthesis of quinazolinone derivatives via multi-component reaction promoted by Pt–MWCNTs nanocomposites.



**Figure 1** Schematic illustration for the generation of Pt NPs supported on MWCNTs.

### 3. Results and discussion

In the present study, we report a green and valid procedure for the synthesis of 2,3-dihydroquinazolin-4(1*H*)-ones via one-pot condensation of isatoic anhydride and aromatic aldehydes with  $\text{NH}_4\text{OAc}$  or aromatic primary amines in the presence of Pt–MWCNTs as a heterogeneous catalyst under ultrasonic irradiation (Scheme 1). The choice of appropriate reaction conditions is important for a successful synthesis. To study the effect of catalyst on the condensation, the one-pot reaction of isatoic anhydride and benzaldehyde with aniline was chosen as a model reaction under ultrasonic irradiation. As shown in Table 1, Pt–MWCNTs were appropriate to catalyze this reaction and a significant improvement was observed in the yield of quinazolinones in the presence of 0.04 g Pt–MWCNTs. Increase in the amount of catalyst did not improve the yield significantly (Table 1, entry 6). In order to improve the yields, we performed reactions using different quantities of reagents. The optimized reactant ratios were found to be 1: 1.2: 1.1: 1 of isatoic anhydride, aniline or ammonium acetate and benzaldehyde in the presence of 0.04 g Pt–MWCNTs at 60 °C for 8 min. In order to investigate the scope and efficiency of this reaction and to study the effect of substituents on the product formation, we applied the optimal protocol for variety of

amines and aldehydes with isatoic anhydride. It is noteworthy that the reaction proceeds efficiently to furnish the corresponding 2,3-dihydroquinazolin-4(1*H*)-ones in excellent yields. It was observed this method had good tolerance for various substitutions. The results are presented in Table 2. It should be noted that the effect of electronic nature and position of substituent on the phenyl rings did not show strong influence on the reaction and the products were obtained in high yields in short reaction times. In our continued study, it was also found that the reaction times are longer and the yields are lower in the absence of ultrasonic irradiation. The results of these reactions are compared under ultrasound irradiation and in an oil

**Table 1** Optimization of amount of catalyst for the three component synthesis of compound 4a.

Entry	Catalyst (g)	Time (min)	Yield (%)
1	0.00	40	20
2	0.01	30	55
3	0.02	25	75
4	0.03	15	85
5	0.04	8	95
6	0.05	8	95

**Table 2** Comparison of time and yields on the formation of compounds **4a–4j** using ultrasound irradiation and conventional heating promoted by Pt–MWCNTs.

Entry	Product	Amine (Ar or NH <sub>4</sub> OAc)	R	Reflux conditions		Ultrasonic irradiation		M.p. (°C)	
				Time (min)	Yield (%)	Time (min)	Yield (%)	Obs.	Lit.
1	<b>4a</b>	Ph	H	35	80	8	95	214–215	214–216 [82]
2	<b>4b</b>	Ph	4-OMe	30	70	12	93	204–206	205–207 [83]
3	<b>4c</b>	4-MeC <sub>6</sub> H <sub>5</sub>	4-F	40	75	15	90	240–242	241–243 [43]
4	<b>4d</b>	Ph	4-Cl	41	71	10	90	223–225	222–224 [84]
5	<b>4e</b>	4-Me C <sub>6</sub> H <sub>5</sub>	4-Me	45	65	20	89	244–246	243–247 [41]
6	<b>4f</b>	4-Cl C <sub>6</sub> H <sub>5</sub>	4-OMe	30	70	15	91	245–247	244–247 [85]
7	<b>4g</b>	5-Cl,2-OH C <sub>6</sub> H <sub>5</sub>	4-Cl	45	65	25	88	235–236	235–237 [41]
8	<b>4h</b>	NH <sub>4</sub> OAc	3-NO <sub>2</sub>	25	85	10	95	196–198	195–196 [84]
9	<b>4i</b>	NH <sub>4</sub> OAc	2-Cl	25	80	12	96	204–205	203–205 [86]
10	<b>4j</b>	NH <sub>4</sub> OAc	4-OH,3-OMe	30	80	20	90	226–228	226–227 [87]


**Table 3** Screening different solvents in the model reaction.

Entry	Solvent	Time (min)	Yield (%)
1	MeOH	15	92
2	PhMe	30	53
3	CH <sub>2</sub> Cl <sub>2</sub>	35	49
4	MeCN	20	85
5	EtOH	8	95
6	THF	40	37


bath and are shown in Table 2. It can be concluded that the synthesis of compounds **4a–4j** is faster and the yields are higher than conventional heating methods. It was apparent that the ultrasound irradiation accelerated this transformation under milder conditions. Its reason may be the phenomenon of cavitation produced by ultrasound waves. Thus the unique combination of Pt–MWCNTs and ultrasound leads to better conversions and a decreased reaction time for this transformation.

For further optimization of the reaction conditions, we then turned our attention to investigate the effect of different solvents on this conversion. The significant rate enhancements were observed when reactions were carried out in ethanol compared with other organic solvents (Table 3). Thus, in order to develop a better reaction system and further optimization, more studies were performed. The possibility of recycling the catalyst was investigated in the model reaction and the activity of the catalyst and yield of the product did not show any significant decrease even after four runs. The results are shown in Fig. 2. In all these cases, the reactions provided a rapid synthetic route, furnishing satisfactorily the respective products that could be isolated and characterized by spectroscopic methods. The structures of all products were characterized on the basis of spectroscopic data particularly <sup>1</sup>H and <sup>13</sup>C NMR spectra.

The representative SEM images of the prepared Pt–MWCNTs catalyst is shown in Fig. 3. It can be observed from SEM images of Pt–MWCNTs that well-dispersed and spherical Pt particles were anchored on the external walls of MWCNTs and the aggregation of alloy nanoparticles was not apparent. Generally, the high dispersion of the Pt nanoparticles on the MWCNTs supports is owing to the existence of uniform surface functional sites on all nanotubes that can be formed in the chemical-wet oxidation of nitric and sulfuric acid.

**Figure 2** Study of recyclability of catalyst in the synthesis of compound **4a**.






**Figure 3** Scanning electron microscopy (SEM) images of Pt-MWCNTs.

The crystalline nature of the Pt nanoparticles was further established by recording the X-ray power diffraction (XRD) pattern of the Pt deposited on MWCNTs sample (Fig. 4). The obvious diffraction peaks at  $2\theta \approx 25.92^\circ$  (002) and

$43.24^\circ$  (100) correspond to the graphitic structure of MWCNTs. The other three peaks  $2\theta = 40.18^\circ$ ,  $46.64^\circ$  and  $67.80^\circ$  can be indexed to the (111), (200) and (220) crystalline planes of Pt NPs, respectively, which indicate the Pt



**Figure 4** XRD patterns of acid treated MWCNTs, Pt nanoparticles and Pt-MWCNTs composite.




Figure 5 FT-IR patterns of (a) acid treated MWCNTs and (b) Pt-MWCNTs composite.





Figure 6 Raman spectra of oxidized MWCNTs and Pt-MWCNTs nanocomposites.

nanoparticles are composed from pure crystalline Pt. According to the Scherrer equation, the particle size of the decorated Pt nanoparticles was about 20–22 nm.

FTIR examination of the acid treated MWCNTs and Pt-MWCNTs catalysts are presented in Fig. 5. The peaks at 3438 and 1726  $\text{cm}^{-1}$  are assigned to stretching vibration of COOH group. The peak at 1627  $\text{cm}^{-1}$  is assigned to the C=C stretching vibration of MWCNTs. The absorption bands at 2919 and 2849  $\text{cm}^{-1}$  are attributed to the stretching vibrations of methylene ( $\text{CH}_2$ ). Stretching modes of Pt-C in

the characteristic frequencies at 1458, 516 and 721  $\text{cm}^{-1}$  can be seen from the FTIR spectrum of Pt-MWCNTs.

The decoration of MWCNTs by Pt nanoparticles is also confirmed by Raman spectra (Fig. 6). The peak at 1307  $\text{cm}^{-1}$  corresponds to amorphous carbon and the defects of pentagons' or heptagons' structure of carbon (D-line) and the peak at 1596  $\text{cm}^{-1}$  indicates the crystalline structure of MWCNT and the  $\text{sp}^2$  hybridized carbon in the graphitic structure (G-line). The intensity ratio of D-band to G-band ( $I_D/I_G$  ratio) is used to study the structural properties and crystallinity of



**Scheme 2** Possible mechanistic route for preparation of quinazolinone derivatives.

MWCNTs. Based on the Raman curves, decrease of the  $I_D/I_G$  ratio from 1.86 in the oxidized MWCNTs to 1.42 in Pt-MWCNTs composites indicates that the restoration of  $sp^2$  structure (reduction of C=C bonds) has occurred in MWCNTs surface and functionalization has been done successfully.

The plausible general mechanistic pathway for this transformation has been outlined in Scheme 2. It is conceivable that the supported metal is coordinated to the oxygen atom of the carbonyl groups in different stages of the reaction activating them for the nucleophilic attack of the amine and amide nitrogen atoms. The reaction can be mechanistically considered to proceed via the initial activation of the isatoic anhydride, then the *N*-nucleophilic amine attack on the carbonyl unit to produce an intermediate **I**, which in turn affords intermediate **II** through decarboxylation reaction. The proton transfer of intermediate **II** affords 2-amino-*N*-substituted-amide **III**. The tautomerization of amide group in intermediate **III** could be catalyzed in the presence of the supported metal. Subsequently, the reaction of activated aldehyde with 2-amino-*N*-substituted-amide **III** proceeds to afford the imine intermediate **IV**. The imine moiety in intermediate **IV** is also activated by catalyst. Cyclization of **IV** to intermediate **V** occurs via intramolecular nucleophilic attack of nitrogen on imine carbon. Then, 1,5-proton shift gives the corresponding products.

#### 4. Conclusions

In conclusion, we applied a newly developed synthetic methodology to synthesize 2,3-dihydroquinazolin-4(1*H*)-ones through the one-pot three-component cyclocondensation of isatoic anhydride, ammonium acetate (or primary aromatic amines)

and aldehydes under ultrasound irradiation. To this end, the focus of our study is the use of Pt-MWCNTs nanocomposites, which are relatively nontoxic and efficient. In addition, in order to obtain the stabilized and high reactive Pt-MWCNTs nanocomposites and to decrease Pt nanoparticle agglomeration in the MWCNTs surface, ultrasound is applied to prepare these nanocomposites. Cavitation can be useful to form stable dispersion of the nanoparticles in the coatings. The present reagent system has not previously been investigated for such condensation. The development of this novel catalyst affords an alternative reagent to prepare these pharmacologically important quinazolinones in good yields. The green procedure, recyclability of catalyst, high yields, short reaction times, simple work-up procedures and use of commercially available starting materials make the present method an interesting approach toward the synthesis of dihydroquinazolinones.

#### Acknowledgment

We are grateful to University of Kashan for financial support (Grant No. 256722/32).

#### References

- [1] A. Strecker, Ueber die künstliche Bildung der Milchsäure und einen neuen, dem Glycocoll homologen Körper, *Liebigs Ann. Chem.* 75 (1850) 27–45.
- [2] B.B. Tour, D.G. Hall, Natural product synthesis using multicomponent reaction strategies, *Chem. Rev.* 109 (2009) 4439–4486.
- [3] J. Zhu, H. Bienayme, *Multicomponent Reactions*, Wiley-VCH, Weinheim, 2005.

- [4] H.A. Parish Jr., R.D. Gilliom, W.P. Purcell, R.K. Browne, R.F. Spirk, H.D. White, Syntheses and diuretic activity of 1,2-dihydro-2-(3-pyridyl)-3H-pyrido[2,3-d]pyrimidin-4-one and related compounds, *J. Med. Chem.* 25 (1982) 98–102.
- [5] M.J. Hour, L.J. Huang, S.C. Kuo, Y. Xia, K. Bastow, Y. Nakanishi, E. Hamel, K.H. Lee, 6-Alkylamino- and 2,3-dihydro-3'-methoxy-2-phenyl-4-quinazolinones and related compounds: their synthesis, cytotoxicity, and inhibition of tubulin polymerization, *J. Med. Chem.* 43 (2000) 4479–4487.
- [6] R.J. Alaimo, H.E. Russell, Antibacterial 2,3-dihydro-2-(5-nitro-2-thienyl) quinazolin-4-(1H)-ones, *J. Med. Chem.* 15 (1972) 335–336.
- [7] D.C. White, T.D. Greenwood, A.L. Downey, J.R. Bloomquist, J.F. Wolfe, Synthesis and anticonvulsant evaluation of some new 2-substituted-3-arylpyrido[2,3-d]pyrimidinones, *Bioorg. Med. Chem.* 12 (2004) 5711–5717.
- [8] X. Cheng, S. Vellalath, R. Goddard, B. List, Direct catalytic asymmetric synthesis of cyclic aminals from aldehydes, *J. Am. Chem. Soc.* 130 (2008) 15786–15787.
- [9] R.J. Abdel-Jalil, W. Volter, M. Saeed, A novel method for the synthesis of 4(3H)-quinazolinones, *Tetrahedron Lett.* 45 (2004) 3475–3476.
- [10] S.E. Lopez, M.E. Rosales, N. Urdaneta, M.V. Godoy, J.E. Charris, The synthesis of substituted 2-aryl 4(3H)-quinazolinones using  $\text{NaHSO}_3/\text{DMA}$ . Steric effect upon the cyclisation-dehydrogenation step, *J. Chem. Res.* 2000 (2000) 258–259.
- [11] V.B. Rao, P. Hanumanthu, C.V. Rantnam, Studies in the formation of heterocyclic rings containing nitrogen, 29: synthesis and thermal-isomerization of 1-alkyl-2-aryl-4(1H)-quinazolinones, *Indian J. Chem.* 18B (1979) 493–496.
- [12] S. John, Quinazolinone derivatives in pharmaceutical research, *Pharmazie* 36 (1981) 583–596.
- [13] S.D. Sharma, V. Kaur, Synthesis of 3-oxa-1-dethiacepham and 3-aza-1-dethiacepham analogs, *Synthesis* (1989) 677–680.
- [14] J. Cizmarik, J. Trupe, Antimicrobial activity of quaternary ammonium-salts of heptacaine, *Pharmazie* 42 (1987) 139–140.
- [15] P.P. Kung, M.D. Casper, K.L. Cook, L. Willson-Lingardo, Structure-activity relationships of novel 2-substituted quinazoline antibacterial agents, *J. Med. Chem.* 42 (1999) 4705–4713.
- [16] J.W. Corbett, S.S. Ko, J.D. Rodgers, L.A. Gearhart, N.A. Magnus, L.T. Bacheler, S. Diamond, S. Jeffey, G.L. Trainor, P.S. Anderson, K. Erickson-Vitanen, Inhibition of clinically relevant mutant variants of HIV-1 by quinazolinone non-nucleoside reverse transcriptase inhibitors, *J. Med. Chem.* 43 (2000) 2019–2030.
- [17] J.F. Liu, J. Lee, A.M. Dalton, G. Bi, L. Yu, C.M. Baldino, E. McElory, M. Brown, Microwave-assisted one-pot synthesis of 2,3-disubstituted 3H-quinazolin-4-ones, *Tetrahedron Lett.* 46 (2005) 1241–1244.
- [18] P. Salehi, M. Dabiri, M. Baghbanzadeh, M. Bahramnejad, One-pot, three-component synthesis of 2,3-dihydro-4(1H)-quinazolinones by Montmorillonite K-10 as an efficient and reusable catalyst, *Synth. Commun.* 36 (2006) 2287–2292.
- [19] S.M. Roopan, F.N. Khan, J.S. Jin, R.S. Kumar, An efficient one pot–three component cyclocondensation in the synthesis of 2-(2-chloroquinolin-3-yl)-2,3-dihydroquinazolin-4(1H)-ones: potential antitumor agents, *Res. Chem. Intermed.* 37 (2011) 919–927.
- [20] P. Salehi, M. Dabiri, M.A. Zolfigol, M. Baghbanzadeh, A novel method for the one-pot three-component synthesis of 2,3-dihydroquinazolin-4(1H)-ones, *Synlett* (2005) 1155–1157.
- [21] M. Dabiri, P. Salehi, M. Baghbanzadeh, M.A. Zolfigol, M. Agheb, S. Heydari, Silica sulfuric acid: an efficient reusable heterogeneous catalyst for the synthesis of 2,3-dihydroquinazolin-4(1H)-ones in water and under solvent-free conditions, *Catal. Commun.* 9 (2008) 785–788.
- [22] L.M. Wang, L. Hu, J.H. Shao, J.J. Yu, L. Zhang, A novel catalyst zinc (II) perfluorooctanoate  $[\text{Zn}(\text{PFO})_2]$ -catalyzed three-component one-pot reaction: Synthesis of quinazolinone derivatives in aqueous micellar media, *J. Fluorine Chem.* 129 (2008) 1139–1145.
- [23] S. Rostamizadeh, A.M. Amani, G.H. Mahdavinia, H. Sepehrian, S. Ebrahimi, Synthesis of some novel 2-aryl-substituted 2,3-dihydroquinazolin-4(1H)-ones under solvent-free conditions using MCM-41- $\text{SO}_3\text{H}$  as a highly efficient sulfonic acid, *Synthesis* (2010) 1356–1360.
- [24] A. Saffar-Teluri, S. Bolouk, One-pot, three-component synthesis of 2,3-dihydroquinazolin-4(1H)-ones using p-toluenesulfonic acid–paraformaldehyde copolymer as an efficient and reusable catalyst, *Monatsh. Chem.* 141 (2010) 1113–1115.
- [25] M.P. Surpur, P.R. Singh, S.B. Patil, S.D. Samant, Expedient one-pot and solvent-free synthesis of dihydroquinazolin-4(1H)-ones in the presence of microwaves, *Synth. Commun.* 37 (2007) 1965–1970.
- [26] J.X. Chen, D.Z. Wu, F. He, M.C. Liu, H.Y. Wu, J.C. Ding, W.K. Su, Gallium (III) triflate-catalyzed one-pot selective synthesis of 2,3-dihydroquinazolin-4(1H)-ones and quinazolin-4(3H)-ones, *Tetrahedron Lett.* 49 (2008) 3814–3818.
- [27] M. Dabiri, P. Salehi, S. Otakesh, M. Baghbanzadeh, G. Kozehgary, A.A. Mohammadi, Efficient synthesis of mono- and disubstituted 2,3-dihydroquinazolin-4(1H)-ones using  $\text{KAl}(\text{SO}_4)_2 \cdot 12\text{H}_2\text{O}$  as a reusable catalyst in water and ethanol, *Tetrahedron Lett.* 46 (2005) 6123–6126.
- [28] M. Dabiri, P. Salehi, M. Bahramnejad, M. Alizadeh, A practical and versatile approach toward a one-pot synthesis of 2,3-disubstituted 4(3H)-quinazolinones, *Monatsh. Chem.* 141 (2010) 877–881.
- [29] S. Rostamizadeh, A.M. Amani, R. Aryan, H.R. Ghaieni, N. Shadjou, Synthesis of new 2-aryl substituted 2,3-dihydroquinazolin-4(1H)-ones under solvent-free conditions, using molecular iodine as a mild and efficient catalyst, *Synth. Commun.* 38 (2008) 3567–3576.
- [30] L.Y. Zeng, C. Cai, Selectively promote the synthesis of mono substituted quinazolin-4(3H)-ones and 2,3-dihydroquinazolin-4(1H)-ones in one-pot, *J. Heterocycl. Chem.* 47 (2010) 1035–1039.
- [31] M. Baghbanzadeh, P. Salehi, M. Dabiri, G. Kozehgary, Water-accelerated synthesis of novel bis-2,3-dihydroquinazolin-4(1H)-one derivatives, *Synthesis* (2006) 344–348.
- [32] H.R. Shaterian, A.R. Oveisi, M. Honarmand, Synthesis of 2,3-dihydroquinazolin-4(1H)-ones, *Synth. Commun.* 40 (2010) 1231–1242.
- [33] K. Niknam, N. Jafarpour, E. Niknam, Silica-bonded N-propylsulfamic acid as a recyclable catalyst for the synthesis of 2,3-dihydroquinazolin-4(1H)-ones, *Chin. Chem. Lett.* 22 (2011) 69–72.
- [34] K.J. Zahed, R. Arjmand, Acetic acid-promoted, efficient, one-pot synthesis of 2, 3-dihydroquinazolin-4(1H)-ones, *Monatsh. Chem.* 142 (2011) 631–635.
- [35] M. Ghashang, K. Azizi, H. Moulavi-Pordanjani, H.R. Shaterian, Eco-friendly and efficient synthesis of 2,3-dihydroquinazolin-4(1H)-ones, *Chin. J. Chem.* 29 (2011) 1617–1623.
- [36] C.L. Yoo, J.C. Fetting, M.J. Kurth, Stannous chloride in alcohol: a one-pot conversion of 2-nitro-N-arylbenzamides to 2,3-Dihydro-1H-quinazolin-4-ones, *J. Org. Chem.* 70 (2005) 6941–6943.
- [37] M. Baghbanzadeh, M. Dabiri, P. Salehi, A new efficient method for the three-component synthesis of 4(3h)-quinazolinones, *Heterocycles* 75 (2008) 2809–2815.
- [38] J.X. Chen, W.K. Su, H.Y. Wu, M.C. Liu, C. Jin, Eco-friendly synthesis of 2,3-dihydroquinazolin-4(1H)-ones in ionic liquids or ionic liquid–water without additional catalyst, *Green Chem.* 9 (2007) 972–975.



- [39] M. Dabiri, P. Salehi, M. Baghbanzadeh, Ionic liquid promoted eco-friendly and efficient synthesis of 2,3-dihydroquinazolin-4(1*H*)-ones, *Monatsh. Chem.* 138 (2007) 1191–1194.
- [40] N.B. Darvatkar, S.V. Bhilare, A.R. Deorukhkar, D.G. Raut, M.M. Salunkhe, [bmim] HSO<sub>4</sub>: an efficient and reusable catalyst for one-pot three-component synthesis of 2,3-dihydro-4(1*H*)-quinazolinones, *Green Chem. Lett. Rev.* 3 (2010) 301–306.
- [41] K. Niknam, M.R. Mohammadzadeh, S. Mirzaee, Silica-bonded S-sulfonic acid as a recyclable catalyst for synthesis of 2,3-dihydroquinazolin-4(1*H*)-ones, *Chin. J. Chem.* 29 (2011) 1417–1422.
- [42] M.Z. Kassaei, S. Rostamizadeh, N. Shadjou, E. Motamedi, M. Esmaelzadeh, An efficient one-pot solvent-free synthesis of 2,3-dihydroquinazolin-4(1*H*)-ones via Al/Al<sub>2</sub>O<sub>3</sub> nanoparticles, *J. Heterocycl. Chem.* 47 (2010) 1421–1424.
- [43] Z.H. Zhang, H.Y. Lu, S.H. Yang, J.W. Gao, Synthesis of 2,3-dihydroquinazolin-4(1*H*)-ones by three-component coupling of isoic anhydride, amines, and aldehydes catalyzed by magnetic Fe<sub>3</sub>O<sub>4</sub> nanoparticles in water, *J. Comb. Chem.* 12 (2010) 643–646.
- [44] Z. Liu, X. Lin, J.Y. Lee, W. Zhang, M. Han, L.M. Gan, Preparation and characterization of platinum-based electrocatalysts on multiwalled carbon nanotubes for proton exchange membrane fuel cells, *Langmuir* 18 (2002) 4054–4060.
- [45] S. Dong, Q. Qiu, Electrodeposition of platinum particles on glassy carbon modified with cobalt porphyrin and nafion film and their electrocatalytic reduction of dioxyge, *J. Electroanal. Chem.* 314 (1991) 223–239.
- [46] T. Frelink, W. Visscher, J.A.R. Van-Veen, Particle size effect of carbon-supported platinum catalysts for the electrooxidation of methanol, *J. Electroanal. Chem.* 382 (1995) 65–72.
- [47] S. Holdcroft, B.L. Funt, Preparation and electrocatalytic properties of conducting films of polypyrrole containing platinum microparticulates, *J. Electroanal. Chem.* 240 (1988) 89–103.
- [48] K. Kinoshita, Particle size effects for oxygen reduction on highly dispersed platinum in acid electrolytes, *J. Electrochem. Soc.* 137 (1990) 845–848.
- [49] K.M. Kost, D.E. Bartak, B. Kazee, T. Kuwana, Electrodeposition of palladium, iridium, ruthenium, and platinum in poly(4-vinylpyridine) films for electrocatalysis, *Anal. Chem.* 62 (1990) 151–157.
- [50] Z. Qi, M.C. Lefebvre, P.G. Pickup, Electron and proton transport in gas diffusion electrodes containing electronically conductive proton-exchange polymers, *J. Electroanal. Chem.* 459 (1998) 9–14.
- [51] S.D. Thompson, L.R. Jordan, M. Forsyth, Platinum electrodeposition for polymer electrolyte membrane fuel cells, *Electrochim. Acta* 46 (2001) 1657–1663.
- [52] C. Wang, M. Waje, X. Wang, J.M. Tang, R.C. Haddon, Y. Yan, Proton exchange membrane fuel cells with carbon nanotube based electrodes, *Nano Lett.* 4 (2004) 345–348.
- [53] K. Yahikozawa, Y. Fujii, Y. Matsuda, K. Nishimura, Y. Takasu, Electrocatalytic properties of ultrafine platinum particles for oxidation of methanol and formic acid in aqueous solutions, *Electrochim. Acta* 36 (1991) 973–978.
- [54] M. Uchida, Y. Aoyama, M. Tanabe, N. Yanagihara, N. Eda, A. Ohta, Influences of both carbon supports and heat-treatment of supported catalyst on electrochemical oxidation of methanol, *J. Electrochem. Soc.* 142 (1995) 2572–2576.
- [55] G. Wu, Y.S. Chen, B.Q. Xu, Remarkable support effect of SWNTs in Pt catalyst for methanol electrooxidation, *Electrochem. Commun.* 7 (2005) 1237–1243.
- [56] R.H. Baughman, A.A. Zakhidov, W.A. Heer, Carbon nanotubes-the route toward applications, *Science* 297 (2002) 787–792.
- [57] T.W. Ebbesen, H.J. Lezec, H. Hiura, J.W. Bennett, H.F. Ghaemi, T. Thio, Electrical conductivity of individual carbon nanotubes, *Nature* 382 (1996) 54–56.
- [58] K.M. Lee, L.C. Li, L.M. Dai, Asymmetric end-functionalization of multi-walled carbon nanotubes, *J. Am. Chem. Soc.* 127 (2005) 4122–4123.
- [59] G. Che, B.B. Lakshmi, C.R. Martin, E.R. Fisher, Metal-nanocluster filled carbon nanotubes: catalytic properties and possible applications in electrochemical energy storage and production, *Langmuir* 15 (1999) 750–758.
- [60] W.Z. Li, C.H. Liang, W.J. Zhou, J.S. Qiu, Z.H. Zhou, G.Q. Sun, Q. Xin, Preparation and characterization of multiwalled carbon nanotube-supported platinum for cathode catalysts of direct methanol fuel cells, *J. Phys. Chem. B* 107 (2003) 6292–6299.
- [61] Y.Y. Shao, G.P. Yin, Y.Z. Gao, P.F. Shi, Durability study of Pt/C and Pt-MWCNTs catalysts under simulated PEM fuel cell conditions, *J. Electrochem. Soc.* 153 (2006) A1093–A1097.
- [62] X. Wang, W.Z. Li, Z.W. Chen, M. Waje, Y.S. Yan, Durability investigation of carbon nanotube as catalyst support for proton exchange membrane fuel cell, *J. Power Sources* 158 (2006) 154–159.
- [63] Y.T. Kim, K. Ohshima, K. Higashimine, T. Uruga, M. Takata, H. Suematsu, T. Mitani, Fine size control of platinum on carbon nanotubes: from single atoms to clusters, *Angew. Chem., Int. Ed.* 45 (2006) 407–411.
- [64] T. Matsumoto, T. Komatsu, H. Nakano, K. Arai, Y. Nagashima, E. Yoo, T. Yamazaki, M. Kijima, H. Shimizu, Y. Takasawa, Efficient usage of highly dispersed Pt on carbon nanotubes for electrode catalysts of polymer electrolyte fuel cells, *Catal. Today* 90 (2004) 277–281.
- [65] N. Rajalakshmi, H. Ryu, M.M. Shaijumon, S. Ramaprabhu, Performance of polymer electrolyte membrane fuel cells with carbon nanotubes as oxygen reduction catalyst support material, *J. Power Sources* 140 (2005) 250–257.
- [66] K.H. Choi, H.S. Kim, T.H. Lee, Electrode fabrication for proton exchange membrane fuel cells by pulse electrodeposition, *J. Power Sources* 75 (1998) 230–235.
- [67] K. Saminathan, V. Kamavaram, V. Veedu, A.M. Kannan, Preparation and evaluation of electrodeposited platinum nanoparticles on in situ carbon nanotubes grown carbon paper for proton exchange membrane fuel cells, *Int. J. Hydrogen Energy* 34 (2009) 3838–3844.
- [68] M.C. Tsai, T.K. Yeh, C.H. Tsai, Methanol oxidation efficiencies on carbon-nanotube-supported platinum and platinum–ruthenium nanoparticles prepared by pulsed electrodeposition, *Int. J. Hydrogen Energy* 36 (2011) 8261–8266.
- [69] X. Li, S. Ge, C.L. Hui, I.M. Hsing, Well-dispersed multiwalled carbon nanotubes supported platinum nanocatalysts for oxygen reduction, *Electrochem. Solid St.* 7 (2004) A286–A289.
- [70] X. Li, I.M. Hsing, The effect of the Pt deposition method and the support on Pt dispersion on carbon nanotubes, *Electrochim. Acta* 51 (2006) 5250–5258.
- [71] Y. Shao, G. Yin, J. Wang, Y. Gao, P. Shi, Multi-walled carbon nanotubes based Pt electrodes prepared with in situ ion exchange method for oxygen reduction, *J. Power Sources* 161 (2006) 47–53.
- [72] J. Wang, G. Yin, Y. Shao, Z. Wang, Y. Gao, Platinum deposition on multiwalled carbon nanotubes by ion-exchange method as electrocatalysts for oxygen reduction, *J. Electrochem. Soc.* 154 (2007) B687–B693.
- [73] T.J. Mason, P. Cintas, in: J. Clark, D. Macquarrie (Eds.), *Handbook of Green Chemistry and Technology*, Blackwell Science, Oxford, 2002.
- [74] T.J. Mason, Sonochemistry and the environment – providing a “green” link between chemistry, physics and engineering, *Ultrason. Sonochem.* 14 (2007) 476–483.
- [75] K.S. Suslick, In *Year Book of Science & the Future Encyclopedia*, Britannica, Chicago, 1994, 138.
- [76] S.D. Lepore, Y. He, Use of sonication for the coupling of sterically hindered substrates in the phenolic Mitsunobu reaction, *J. Org. Chem.* 68 (2003) 8261–8263.

- [77] J.S. Yadav, B.V.S. Reddy, K.B. Reddy, K.S. Raj, A.R. Prasad, Ultrasound-accelerated synthesis of 3,4-dihydropyrimidin-2(1H)-ones with ceric ammonium nitrate, *J. Chem. Soc., Perkin Trans. 1* (2001) 1939–1941.
- [78] S. Streiff, N. Ribeiro, L. Desaubry, Unexpected cleavage of tetrahydrofuran by catalytic reductive lithiation, *Chem. Commun.* (2004) 346–347.
- [79] A. Gedanken, Using sonochemistry for the fabrication of nanomaterials, *Ultrason. Sonochem.* 11 (2004) 47–55.
- [80] K.S. Suslick, M.M. Fang, T. Hyeon, M.M. Mdleleni, Applications of sonochemistry to materials synthesis, *Sonochem. Sonolumin.* (1999) 291–320.
- [81] O.Y. Ivanshina, M.E. Tamm, E.V. Gerasimova, M.P. Kochugaeva, M.N. Kirikova, S.V. Savilov, L.V. Yashina, Synthesis and electrocatalytic activity of platinum nanoparticle/carbon nanotube composites, *Inorg. Mater.* 47 (2011) 618–625.
- [82] Z. Song, X. Wan, S. Zhao, A novel catalyst cobalt m-nitrobenzenesulfonate-catalyzed highly efficient synthesis of substituted-quinazolin-4-(1H)-ones, *Indian J. Chem. Technol.* 19 (2012) 118–123.
- [83] M. Wang, T.T. Zhang, J.J. Gao, Y. Liang, Cation-exchange resin as an efficient heterogeneous catalyst for one-pot three-component synthesis of 2,3-dihydro-4(1H)-quinazolinones, *Chem. Heterocycl. Compd.* 48 (2012) 897–902.
- [84] M. Wang, T.T. Zhang, Y. Liang, J.J. Gao, Efficient synthesis of mono- and disubstituted 2,3-dihydroquinazolin-4(1H)-ones using copper benzenesulfonate as a reusable catalyst in aqueous solution, *Monatsh. Chem.* 143 (2012) 835–839.
- [85] Q.Y. Zhuang, Y.C. Fu, D. Tang, Y. Zha, L. Rong, S. Tu, Synthesis of 2,3-diaryl-2,3-dihydroquinazolin-4(1H)-one under solvent-free conditions, *Chin. J. Org. Chem.* 30 (2010) 1405–1409.
- [86] Z. Song, L. Liu, Y. Wang, X. Sun, Efficient synthesis of mono- and disubstituted 2,3-dihydroquinazolin-4(1H)-ones using aluminum methanesulfonate as a reusable catalyst, *Res. Chem. Intermed.* 38 (2012) 1091–1099.
- [87] M. Wang, T.T. Zhang, Z.G. Song, Eco-friendly synthesis of 2-substituted-2,3-dihydro-4(1H)-quinazolinones in water, *Chin. Chem. Lett.* 22 (2011) 427–430.